

Ingham type theorems and multiplier method for some specific domains

Michel Mehrenberger

IRMA, Université de Strasbourg

mehrenbe@math.unistra.fr

Abstract: Ingham type theorems and the multiplier method are complementary tools for dealing with the observability of wave type equations. We give here some specific examples and try to compare and make the link between the different approaches. This work is a collaboration with Vilmos Komornik and Paola Loreti.

- [1.] A. Barhoumi, V. Komornik, M. Mehrenberger, A vectorial Ingham-Beurling type theorem, special issue of the journal *Annales Univ. Sci. Budapest*, dedicated to Professor Laszlo Simon on the occasion of his 70th birthday, Vol 53 (2010), 17–32.
- [2.] P. Loreti, M. Mehrenberger, Observabilité uniforme de l'équation des ondes 1D [Uniform observability of the 1D wave equation], Paris-Sud Working Group on Modelling and Scientific Computing 2007–2008, 68–79, *ESAIM Proc.*, 25, EDP Sci., Les Ulis, 2008.
- [3.] P. Loreti, M. Mehrenberger, An Ingham type proof for a two-grid observability theorem, *ESAIM Control Optim. Calc. Var.* 14 (2008), no. 3, 604–631.
- [4.] M. Mehrenberger, An Ingham type proof for the boundary observability of a N-d wave equation, *C. R. Math. Acad. Sci. Paris* 347 (2009), no. 1-2, 63–68.